

GAME COUNTS IN NORTH-WEST NAMIBIA

Hobatere concession

June 2018


Total Population Estimates

Species	Population estimate	Lower 95% CL	Upper 95% CL
Gemsbok (HN)	80	10	510
Kudu (HN)	-	-	-
Springbok (HN)	520	100	2595
Hartmann's Zebra (HN)	340	160	730


All above estimates are derived using DISTANCE analysis. This takes account of drop off in detection with distance from the transect line. Model selection: U = uniform key; HN = half normal

Total number of animals seen each year

Species	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2015	2016	2017	2018
Baboon				4											1	20
Eland	52	19		12	10	12	9						45	5		
Elephant	6	2	1				2	3			20	1	1	3	10	
Gemsbok	83	217	251	108	158	117	37	52	80	82	40	90	24	15	48	18
Giraffe	54	47	15	63	31	24	18	35	11	1	12	9	29	33	26	13
Jackal		9	3		5	4	5	3	3	8		3	8	7	4	2
Kudu	17	7	4	11	15	15	5	3	3	1		3	1	14	6	
Ostrich	3	2	4		1	4									1	
Springbok	130	66	170	154	194	52	51	65	50	94	51	60	34	53	38	62
Steenbok	2	2	1	3	3	1	4				1	3	2	5	1	1
Warthog	3	9	8	1	10	5	1		2			1	4	1	3	
H. Zebra	190	148	53	83	90	147	63	77	105	64	65	27	129	130	52	80



Trends - Number of animals per 100km




Animals seen during this count and minimum estimates

Values without brackets are numbers of animals seen along transects. Values inside brackets are minimum estimates assuming all animals within 500m on each side of the transect line are detected i.e. there is no adjustment for drop off in detection with distance from the transect line.


These values are significantly lower than the totals indicated in the top left table as the total estimates take account of species detection curves.

Total Route km	62
Total area (km ²)	258
Number of routes	3
% area excluded	5

Species	Count	Minimum Estimate
Gemsbok	18	(37)
Giraffe	13	(44)
Kudu		
Ostrich		
Springbok	62	(174)
Steenbok	1	(2)
H. Zebra	80	(246)





Rainfall



The rainfall season is from July to June and values are an average for the whole area. The year represents the season immediately prior to the count.

Vegetation



NDVI is a measure of the density of chlorophyll in vegetation cover. It can be used as an indicator of the amount of biomass available to wildlife. The map shows the NDVI status in the current year (Mar-May) and the trend indicates the average deviation from the long-term mean (2001-2018).