

# GAME COUNTS IN NORTH-WEST NAMIBIA

## Conservancies north of the veterinary fence

May 2019

### Total Population Estimates


Species	Population estimate	Lower 95% CL	Upper 95% CL
Gemsbok (U)	3,820	2,290	6,380
Kudu (HN)	730	360	1,490
Ostrich (U)	5,070	3,680	6,980
Springbok (HN)	39,950	27,900	57,210
Steenbok (HN)	3,920	2,500	6,150
Hartmann's Zebra (HN)	3,710	2,290	6,020

All above estimates are derived using DISTANCE analysis. This takes account of drop off in detection with distance from the transect line. They are conservative estimates as 35% of the count area is not sampled (due to inaccessibility) and is consequently assumed to hold no animals. Model selection: U = uniform key; HN = half normal

### Total number of animals seen each year

Species	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Baboon	45	44	55	73	253	61	234	105	105	125	17	136	272	123	235	331	132	114	292
Cheetah	2	1	7	2	10	1	1	4	3	1		3	2	1			3	1	2
Duiker	4	1	2	1			6			5	4	1	3	6	3	3		2	6
Eland	11						36	5	30		13	2		2		16	5	9	1
Elephant	16	10	24	5	31	1	4	8	20	10	1	9	8	4	12	14	18	11	31
Gemsbok	809	975	2,152	1,348	1,845	1,508	2,649	1,337	1,580	1,493	1,257	1,911	1,401	1,101	575	524	586	626	491
Giraffe	64	89	75	111	59	140	99	85	86	163	182	252	186	152	178	256	209	219	337
Hyaena	1						2						6	1		1		1	
Jackal	11	26	25	45	28	42	50	19	35	31	28	26	41	35	24	21	35	18	10
Klipspringer	3	5	6	9		3	5		15	3	2	8	4	11		5		2	
Kudu	20	30	18	46	61	43	54	10	88	71	23	56	84	86	63	100	35	18	41
Ostrich	298	394	454	427	458	469	534	339	702	454	451	439	555	502	422	496	435	398	517
Springbok	5,078	5,823	4,692	3,748	6,865	4,527	6,939	13,999	4,342	6,569	6,136	4,463	4,640	3,662	3,210	6,015	5,028	4,816	2,786
Steenbok	20	32	41	115	64	53	80	34	54	60	44	90	113	93	97	96	61	35	68
Warthog		5		2	1	2	6	2		3	2	5	10	1		3	2	3	
H. Zebra	283	189	302	367	568	576	558	585	950	1,271	970	748	619	1,028	256	993	1,177	1,529	382

### Trends - Number of animals per 100km



### Animals seen during this count and minimum estimates


	Total	Anabeb	Ehi-Rovipuka	Marientluss	Okangundumba	Okondjombo	Omatendeka	Ombujokanguindi	Ongongo	Orupembe	Orupupa	Otjambangu	Ojikondavirongo	Otju-west	Otuzemba	Ozondundu	Puros	Sanitatas	Sesfontein
Total Route km	3,247	233	273	281	139	192	211	158	64	292	159	56	135	88	102	77	328	151	307
Total area (km2)	28,288	1,636	1,979	3,034	1,130	1,643	1,613	657	619	2,616	1,775	348	1,067	1,208	741	743	3,564	1,446	2,469
Number of routes	74	5	5	6	3	4	4	3	5	6	3	1	4	2	3	2	6	4	8
% area excluded	35	51	28	28	29	16	48	26	53	10	44	74	58	71	26	55	31	28	42

Species	Total	Anabeb	Ehi-Rovipuka	Marientluss	Okangundumba	Okondjombo	Omatendeka	Ombujokanguindi	Ongongo	Orupembe	Orupupa	Otjambangu	Ojikondavirongo	Otju-west	Otuzemba	Ozondundu	Puros	Sanitatas	Sesfontein
Gemsbok	7		87 (837)	5 (40)	1 (4)		197 (1,549)								70 (354)	78 (736)	53 (213)		
Giraffe	21 (42)	110 (372)	3 (6)	7 (14)	125 (250)			6 (12)	3 (18)							22 (44)	23 (46)	17 (34)	
Kudu	5 (24)	7 (26)	1 (7)		1 (4)			4 (23)	2 (13)						3 (17)	14 (40)		4 (22)	
Ostrich	9 (19)	10 (65)	64 (263)	8 (44)	8 (45)	26 (105)	45 (183)	21 (106)	81 (639)	26 (156)	25 (40)	2 (7)	35 (138)	1 (3)	7 (31)	54 (117)	19 (106)	76 (233)	
Springbok	338 (653)	104 (384)	143 (900)	57 (317)	773 (3,498)	122 (486)	131 (590)	7 (35)	233 (1,756)	42 (270)	156 (248)	36 (120)	1 (4)		187 (923)	56 (77)	45 (205)	355 (1,581)	
Steenbok		1 (9)	1 (7)	4 (22)	3 (23)	5 (20)	9 (39)	1 (4)	9 (79)	18 (116)	3 (5)	1 (3)		3 (17)	2 (6)		4 (22)	4 (9)	
Hartmann's Zebra	52 (121)	49 (144)	21 (152)		37 (137)				69 (580)						4 (11)	56 (323)	6 (42)	88 (451)	


Values without brackets are numbers of animals seen along transects. Values inside brackets are minimum estimates assuming all animals within 500m on each side of the transect line are detected i.e. there is no adjustment for drop off in detection with distance from the transect line. In addition, for springbok, gemsbok and giraffe, large groups were excluded from extrapolations and added afterwards.

The sum of these values will be significantly lower than the totals indicated in the top left table as the total estimates take account of species detection curves.


### Rainfall



### Vegetation



### Mortalities



The rainfall season is from July to June and values are an average for the whole area. The year represents the season immediately prior to the count.

NDVI is a measure of the density of chlorophyll in vegetation cover. It can be used as an indicator of the amount of biomass available to wildlife. The map shows the NDVI status in the current year (Mar-May) and the trend indicates the average deviation from the long-term mean (2001-2018).

Total number of carcasses from all species recorded in the Event Book